

Asymmetric Unimodal Maps: Some Results from q -generalized Bit Cumulants

Uğur Tirnaklı*

*Centro Brasileiro de Pesquisas Físicas, Rua Xavier Sigaud 150, 22290-180 Rio de Janeiro - RJ, Brazil
and*

Department of Physics, Faculty of Science, Ege University, 35100 Izmir-Turkey

In this study, using q -generalized bit cumulants (q is the nonextensivity parameter of the recently introduced Tsallis statistics), we investigate the asymmetric unimodal maps $x_{t+1} = 1 - a|x_t|^{z_i}$ ($i = 1, 2$ correspond to $x_t > 0$ and $x_t < 0$ respectively, $z_i > 1$, $0 < a \leq 2$, $t = 0, 1, 2, \dots$). The study of the q -generalized second cumulant of these maps allows us to determine, for the first time in the literature, the dependence of the inflexion parameter pairs (z_1, z_2) to the nonextensivity parameter q . This behaviour is found to be very similar to that of the logistic-like maps ($z_1 = z_2 = z$) reported recently by Costa et al. [Phys. Rev. E 56 (1997) 245].

PACS Number(s): 05.45.-a, 05.20.-y, 05.70.Ce

In recent years, sensitivity to initial conditions of nonlinear dynamical systems has started to be studied with an increasing interest. Among them, dissipative systems of low-dimensional maps [1–4], self-organized criticality [5], symbolic sequences [6] and conservative systems of long-ranged many-body Hamiltonians [7,8], low-dimensional maps [9] can be enumerated. We focus here on the one-dimensional dissipative maps. As it has already been well studied in the literature, to investigate the sensitivity to initial conditions of any kind of one-dimensional maps on the onset to chaos, it is possible to introduce

$$\xi(t) = \lim_{\Delta x(t) \rightarrow 0} \frac{\Delta x(t)}{\Delta x(0)} \quad , \quad (1)$$

(where $\Delta x(0)$ and $\Delta x(t)$ are discrepancies of the initial conditions at times 0 and t) which, if the Lyapunov exponent $\lambda_1 \neq 0$, satisfies the differential equation $d\xi/dt = \lambda_1 \xi$, thus

$$\xi(t) = e^{\lambda_1 t} \quad , \quad (2)$$

whereas, if $\lambda_1 = 0$, it satisfies $d\xi/dt = \lambda_q \xi^q$, thus

$$\xi(t) = [1 + (1 - q)\lambda_q t]^{\frac{1}{1-q}} \quad (q \in \mathcal{R}) \quad (3)$$

which recovers the standard case (namely Eq.(2)) for $q = 1$. Here, q is the nonextensivity parameter of recently introduced Tsallis statistics [10] and for $q \neq 1$ case, it is evident that Eq.(3) yields a *power-law* sensitivity to initial conditions. At the onset of chaos, $\xi(t)$ presents strong fluctuations reflecting the fractal-like structure of the critical attractor and Eq.(3) delimits the power-law growth of the upper bounds of $\xi(t)$. These upper bounds ($\xi \propto t^{\frac{1}{1-q}}$) allow us to calculate the values of q for the map under consideration. This method of finding numerical values of q has been successfully used for the logistic map [1], a family of logistic-like maps [2], the circle map [3] and a family of circular-like maps [4]. Beside this method, another one has been developed by Lyra and Tsallis [3] by looking at the geometrical aspects of dynamical attractors at the threshold to chaos. Using the multifractal singularity spectrum $f(\alpha)$ [11], they proposed the scaling relation

$$\frac{1}{1-q} = \frac{1}{\alpha_{min}} - \frac{1}{\alpha_{max}} \quad (4)$$

*e-mail: tirnakli@sci.ege.edu.tr

where α_{max} (α_{min}) is the most rarefied (concentrated) region of the multifractal singularity spectrum of the attractor. This relation presents the second method of calculating the q values once the scaling properties of the dynamical attractor are known. It has already been shown that for all above-mentioned one-dimensional dissipative maps, the values of q index calculated within these two different methods are the same (within a good precision) for any given value of the inflexion parameter z of the used map. In addition to this, another important information which comes from all these works is the determination of the behaviour of the q index as a function of the inflexion parameter z . As it can be seen from Fig.2 of [2] and from the Table of [4], it seems that when $z \rightarrow \infty$, q approaches 1.

The purpose of this paper is to show numerically whether such kind of behaviour is verified for asymmetric unimodal maps (AUMs)

$$x_{t+1} = \begin{cases} 1 - a|x_t|^{z_1} & \text{if } x_t > 0 \\ 1 - a|x_t|^{z_2} & \text{if } x_t < 0 \end{cases} \quad (5)$$

where $z_{1,2} > 1$, $0 < a \leq 2$ and $t = 0, 1, 2, \dots$. Unfortunately, up to now, for the AUMs, neither the first nor the second method can yield satisfactory results for the prediction of q values of any inflexion parameter pair (z_1, z_2) [12]. The problem of the first method might be originated from the prediction of the critical a_c values at the chaos threshold with enough precision, whereas the problem of the second method might be related to the numerical procedure used to estimate the $f(\alpha)$ curve, whose most rarefied region (α_{max}) is usually poorly sampled. Another fact which yields this failure might be the nonuniversal behaviour of AUMs reported in [13,14] (for example, it is shown that AUMs fail to exhibit the metric universality of Feigenbaum). Since the values of q are not available for AUMs, clearly it is not possible to see the behaviour of q as a function of (z_1, z_2) pairs by using the above-mentioned two methods. In this paper, in order to see this behaviour, *without* finding the precise values of q for (z_1, z_2) pairs, we use another technique based on the very recent generalization of bit cumulants for chaotic systems within Tsallis statistics [15,16]. To explain this technique, let us recall the main results of [15,16]. The generalized second cumulant (or heat capacity) is given by

$$C_2^{(q)} = \frac{q}{(q-1)^2} \left[\langle \rho^{2q-1} \rangle - \langle \rho^q \rangle^2 \right] \quad (6)$$

where ρ is the natural invariant density [17]. As $q \rightarrow 1$, we have

$$C_2^{(1)} = \langle (\ln \rho)^2 \rangle - \langle \ln \rho \rangle^2 \quad (7)$$

which is equivalent to the standard definition of the second cumulant [17,18]. It is shown [15,16] that there is a kind of scaling between $C_2^{(1)}$ and $C_2^{(q)}$ which is evident from a $C_2^{(1)}$ vs $C_2^{(q)}$ plot, where it is easily seen that most of the points can be fitted to a straight line. The slope of this line gives the scaling factor between $C_2^{(1)}$ and $C_2^{(q)}$ and also provides us a tool to fulfill our aim in this paper. In Fig.1, we illustrate this slope plotting $C_2^{(1)}$ vs $C_2^{(q)}$ curve for a representative (z_1, z_2) pair. It is evident that most of the data points fall onto a straight line which yields a well-defined slope (in fact, we encountered that a few points deviate from this straight line especially when the values of the inflexion parameters start to be very different from the standard case (namely, $z_1 = z_2$), but since such data points are very few, we prefer to calculate the slope using a linear regression –with an error of ± 0.01 for each estimation of the slope– to data points which fall onto this straight line). As it is evident from Fig.2, when $q \rightarrow 1$, naturally $C_2^{(q)} \rightarrow C_2^{(1)}$ and thus the slope also tends to unity. Therefore, now we have another parameter (which behaves exactly the same as q) from where we can check the behaviour of the q index as a function of (z_1, z_2) pairs *without knowing the exact values of q !* Calculating the above-mentioned slope for various values of (z_1, z_2) pairs, we are able to estimate the behaviour of q index as a function of inflexion parameters without finding any q value. Fig.3 represents this behaviour for $(2, z_2), (z_1, 2), (3, z_2)$ and $(z_1, 3)$ cases. It seems from the figure that as $z_2 - z_1 \rightarrow \pm\infty$, the above-mentioned slope (and thus, q index) *approaches to unity!* This tendency is exactly the same as that observed for a family of logistic-like maps [2] and a family of circular-like maps [4], namely, as $z \rightarrow \infty$, q approaches 1.

Summing up, for the first time (to the best of our knowledge), we manage to study the asymmetric unimodal maps in such a way that it became possible to relate it with the nonextensivity parameter q of Tsallis statistics. We were able to show that the dependence of the inflexion parameter pairs (z_1, z_2) of these maps to the q index is very similar to that observed for other one-dimensional maps reported so

far. Although the determination of the exact values of q index for (z_1, z_2) pairs is still lacking, we hope that present work would be considered as a first attempt on this line and will accelerate other studies which are, no doubt, highly welcome.

ACKNOWLEDGMENTS

I acknowledge the partial support of BAYG-C program of TUBITAK (Turkish agency) and CNPq (Brazilian agency) as well as the support from Ege University Research Fund under the project number 98FEN025.

-
- [1] C. Tsallis, A.R. Plastino and W.-M. Zheng, *Chaos, Solitons and Fractals* **8** (1997) 885.
 - [2] U.M.S. Costa, M.L. Lyra, A.R. Plastino and C. Tsallis, *Phys. Rev. E* **56** (1997) 245.
 - [3] M.L. Lyra and C. Tsallis, *Phys. Rev. Lett.* **80** (1998) 53.
 - [4] U. Tirnakli, C. Tsallis and M.L. Lyra, *Eur. Phys. J. B* **11** (1999) 309.
 - [5] F.A. Tamarit, S.A. Cannas and C. Tsallis, *Eur. Phys. J. B* **1** (1998) 545; A.R.R. Papa and C. Tsallis, *Phys. Rev. E* **57** (1998) 3923.
 - [6] M. Buiatti, P. Grigolini and L. Palatella, *Physica A* **268** (1999) 214.
 - [7] C. Anteneodo and C. Tsallis, *Phys. Rev. Lett.* **80** (1998) 5313.
 - [8] V. Latora, A. Rapisarda and S. Ruffo, *Phys. Rev. Lett.* **80** (1998) 692; A. Torcini and M. Antoni, *Phys. Rev. E* **59** (1999) 2746.
 - [9] V. Latora and M. Baranger, *Phys. Rev. Lett.* **82** (1999) 520.
 - [10] C. Tsallis, *J. Stat. Phys.* **52** (1988) 479; E.M.F. Curado and C. Tsallis, *J. Phys. A* **24** (1991) L69 [corrigenda: **24** (1991) 3187 and **25** (1992) 1019].
 - [11] T.C. Halsey et al., *Phys. Rev. A* **33** (1986) 1141.
 - [12] U. Tirnakli, C. Tsallis and M.L. Lyra, unpublished results.
 - [13] R.V. Jensen and L.K.H. Ma, *Phys. Rev. A* **31** (1985) 3993.
 - [14] M.C. de Sousa Vieira, E. Lazo and C. Tsallis, *Phys. Rev. A* **35** (1987) 945.
 - [15] R.S. Johal and R. Rai, '*Nonextensive thermodynamic formalism for chaotic dynamical systems*', cond-mat/9909218.
 - [16] R. Rai and R.S. Johal, '*Generalized bit cumulants for chaotic systems: Numerical results*', cond-mat/9909288.
 - [17] F. Schlogl and E. Scholl, *Z. Phys. B* **72** (1988) 231.
 - [18] C. Beck and F. Schlogl, *Thermodynamics of chaotic systems: an introduction* (Cambridge University Press, Cambridge, 1993).

Figure Captions

Figure 1 : The scaling between the standard second cumulant $C_2^{(1)}$ and the generalized second cumulant $C_2^{(q)}$ for a representative value of (z_1, z_2) pairs.

Figure 2 : The behaviour of the slope as a function of q index for a representative value of (z_1, z_2) pairs.

Figure 3 : The behaviour of the slope as a function of $z_2 - z_1$ values for a family of four different (z_1, z_2) pairs.

Fig 1




Fig 2




Fig 3

